

Zero-one laws: A quantitative insight

Alex Wan
(jww Thomas Powell)

Department of Computer Science, University of Bath



'Every mathematician believes that he is ahead of the others. The reason none state this belief in public is because they are intelligent people.' - Andrey Kolmogorov

Zero-one laws are theorems which state that events of a certain type occur with probability 0 or 1.

Such events include tail events. A tail event T is associated with a sequence of independent events (A_n) . By changing a finite number of occurrences concerning the events A_n , it doesn't change the occurrence of the event T .

For an analogy, one can think of a sequence of real numbers converging; the tail of the sequence dictates the behaviour.

Borel's zero-one law

Given a sequence of events (A_n) , the event A_∞ is the event that infinitely many of the A_n occur. A_∞ is a tail event.

$$A_\infty = \lim_{k \rightarrow \infty} \bigcup_{i=k}^{\infty} A_i = \bigcap_{k=1}^{\infty} \bigcup_{i=k}^{\infty} A_i \equiv \forall k \exists i \geq k A_i.$$

Theorem (Borel's zero-one law: Qualitative version)

Let (A_n) be a sequence of independent events; then $\mathbb{P}(A_\infty)$ can only assume only the values zero or one.

Borel's zero-one law in logical form

$$[\mathbb{P}(A_\infty) = 0 \vee \mathbb{P}(A_\infty) = 1]$$

$$\Leftrightarrow \neg [\mathbb{P}(A_\infty) \in (0, 1)]$$

$$\Leftrightarrow \neg [\mathbb{P}(A_\infty) > 0 \wedge \mathbb{P}(A_\infty) < 1]$$

$$\Leftrightarrow \neg \left[\lim_{k \rightarrow \infty} \mathbb{P}(\bigcup_{i=k}^{\infty} A_i) > 0 \wedge \lim_{k \rightarrow \infty} \mathbb{P}(\bigcup_{i=k}^{\infty} A_i) < 1 \right]$$

$$\Leftrightarrow \neg \left[\exists \varepsilon > 0 \forall l \in \mathbb{N} \mathbb{P}(\bigcup_{i=l}^{\infty} A_i) \geq \varepsilon \right. \\ \left. \wedge \exists \delta > 0 \exists k \in \mathbb{N} \mathbb{P}(\bigcup_{i=k}^{\infty} A_i) \leq 1 - \delta \right]$$

$$\Leftrightarrow \neg \left[\exists \varepsilon > 0 \forall l \in \mathbb{N} \exists n \in \mathbb{N} \mathbb{P}(\bigcup_{i=l}^n A_i) \geq \varepsilon \right. \\ \left. \wedge \exists \delta > 0 \exists k \in \mathbb{N} \forall m \in \mathbb{N} \mathbb{P}(\bigcup_{i=k}^m A_i) \leq 1 - \delta \right]$$

Borel's zero-one law in metastable form

$$\Leftrightarrow \neg \left[\exists \varepsilon > 0 \forall l \in \mathbb{N} \exists n \in \mathbb{N} \mathbb{P} \left(\bigcup_{i=l}^n A_i \right) \geq \varepsilon \right. \\ \left. \wedge \exists \delta > 0 \exists k \in \mathbb{N} \forall m \in \mathbb{N} \mathbb{P} \left(\bigcup_{i=k}^m A_i \right) \leq 1 - \delta \right]$$

$$\Leftrightarrow \neg \left[\exists \varepsilon > 0 \exists g : \mathbb{N} \rightarrow \mathbb{N} \forall l \in \mathbb{N} \mathbb{P} \left(\bigcup_{i=l}^{g(l)} A_i \right) \geq \varepsilon \right. \\ \left. \wedge \exists \delta > 0 \exists k \in \mathbb{N} \forall m \in \mathbb{N} \mathbb{P} \left(\bigcup_{i=k}^m A_i \right) \leq 1 - \delta \right]$$

$$\Leftrightarrow \neg \left[\exists \delta, \varepsilon > 0 \exists k \in \mathbb{N} \exists g : \mathbb{N} \rightarrow \mathbb{N} \forall l, m \in \mathbb{N} \right. \\ \left. \mathbb{P} \left(\bigcup_{i=l}^{g(l)} A_i \right) \geq \varepsilon \wedge \mathbb{P} \left(\bigcup_{i=k}^m A_i \right) \leq 1 - \delta \right]$$

$$\Leftrightarrow \forall \delta, \varepsilon > 0 \forall k \in \mathbb{N} \forall g : \mathbb{N} \rightarrow \mathbb{N} \exists l, m \in \mathbb{N} \\ \left[\mathbb{P} \left(\bigcup_{i=l}^{g(l)} A_i \right) < \varepsilon \vee \mathbb{P} \left(\bigcup_{i=k}^m A_i \right) > 1 - \delta \right]$$

Borel's zero-one law in a quantitative statement

Theorem (Borel's zero-one law: Quantitative version)

Let (A_n) be a sequence of independent events. Then for all $\delta, \varepsilon \in (0, 1)$, k a natural number and $g : \mathbb{N} \rightarrow \mathbb{N}$ with $g(r) \geq r$ for all $r \in \mathbb{N}$, we have

$$\exists l \leq h^{(\alpha)}(k) \left[\mathbb{P} \left(\bigcup_{i=l}^{g(l)} A_i \right) < \varepsilon \vee \mathbb{P} \left(\bigcup_{i=k}^m A_i \right) > 1 - \delta \right],$$

where

$$\begin{aligned} \alpha &:= \left\lceil \log\left(\frac{1}{\delta}\right) / \varepsilon \right\rceil + 1, \\ h(n) &:= g(n) + 1, \\ m &:= g \circ h^{(\alpha)}(k). \end{aligned}$$

A generalization of Borel's zero-one law

Arthan and Oliva studied a generalized version of the Borel-Cantelli lemmas, namely the Erdős-Rényi theorem in [1].

Similarly, we can give a generalization of Borel's zero-one law:

Theorem

Let (A_n) be a sequence of pairwise independent events. Then for all $\delta, \varepsilon \in (0, 1)$, k natural number and $g : \mathbb{N} \rightarrow \mathbb{N}$ with $g(r) \geq r$ for all $r \in \mathbb{N}$,

$$\exists l \leq h^{(\beta(k))}(k) \left[\mathbb{P} \left(\bigcup_{i=l}^{g(l)} A_i \right) < \varepsilon \vee \mathbb{P} \left(\bigcup_{i=k}^m A_i \right) > 1 - \delta \right],$$

where

$$\begin{aligned} \beta(k) &:= \left\lfloor \frac{2k}{\varepsilon} \right\rfloor \\ h(n) &:= g(n) + 1, \\ m &:= g \circ h^{(\beta(k))}(k). \end{aligned}$$

Moving away from Borel's zero-one law

Tail events of the form A_∞ can be presented in a more abstract way:

$$T := \bigcap_{n=0}^{\infty} \bigcup_{k=n}^{\infty} B(n, k) \equiv \forall n \exists k \geq n B(n, k),$$

where the events $B(n, k)$ satisfies the following properties:

- 1 if $k < n$, then $B(n, k) = \emptyset$;
- 2 $B(n, m) \cup B(l, k) \subseteq B(n, k)$ for all $n \leq m < l \leq k$;
- 3 $B(n, m)$ and $B(l, k)$ are independent for all $n \leq m < l \leq k$.

We take $B(n, k) = \bigcup_{i=k}^n A_i$ to recover the event A_∞ .

Theorem (A more generalized quantified zero-one law)

Let the abstract event $B(n, k)$ satisfy the previous properties. Then for all $\delta, \varepsilon \in (0, 1)$, k a natural number and $g : \mathbb{N} \rightarrow \mathbb{N}$ with $g(r) \geq r$ for all $r \in \mathbb{N}$,

$$\exists l \leq h^{(\alpha)}(k) [\mathbb{P}(B(l, g(l))) < \varepsilon \vee \mathbb{P}(B(k, m)) > 1 - \delta.],$$

where

$$\begin{aligned}\alpha &:= \left\lceil \log\left(\frac{1}{\delta}\right) / \varepsilon \right\rceil + 1, \\ h(n) &:= g(n) + 1, \\ m &:= g \circ h^{(\alpha)}(k).\end{aligned}$$

Examples

Examples where the framework may apply:

- **Rademacher sequences:** Given a sequence of real numbers (a_n) and a sequence of independent and identically distributed random variables (ϵ_n) taking values in $\{-1, 1\}$, does the sum

$$\sum_{n=1}^{\infty} \epsilon_n a_n$$


almost surely converge?

- **Bond percolation:** Given a random process on the edges of a lattice, almost surely does there exist an infinitely connected component?

Percolation: N-squares

Let $B(n, k)$ be the statement

'there exists no filled path from the n -square to k -square'.



Theorem (Kolmogorov's zero-one law)

Let (A_n) be a sequence of independent events and denote by $\mathcal{B}^{(n)} = \mathcal{B}(A_{n+1}, A_{n+2}, \dots)$ the smallest σ -algebra which contains all the sets A_k with subscript $k \geq n + 1$. Suppose that T is an event which belongs to $\mathcal{B}^{(n)}$ for all n , then either $\mathbb{P}(T) = 0$ or $\mathbb{P}(T) = 1$.

Summary

For tail events, we can finitize the associated zero-one law statement and give a uniform bound (independent of the sequence of the event) to say probabilistically if we are close to one case or the other.

Not only are finitizations interesting in their own right, but they can be used to extract numerical rates from existential proofs that use 0-1 laws as a lemma.

From this, it is natural to finitize more complicated zero-one laws, such as Kolmogorov's zero-one law and Hewitt-Savage's zero-one law.

Thank you!

References

- [1] Rob Arthan and Paulo Oliva. “On the Borel-Cantelli Lemmas, the Erdős-Rényi Theorem, and the Kochen-Stone Theorem”. In: *Journal of Logic and Analysis* 13 (Dec. 2021). ISSN: 1759-9008. DOI: [10.4115/jla.2021.13.6](https://doi.org/10.4115/jla.2021.13.6).
- [2] Geoffrey Grimmett. *Percolation*. Springer Berlin Heidelberg, 1999. ISBN: 9783662039816. DOI: [10.1007/978-3-662-03981-6](https://doi.org/10.1007/978-3-662-03981-6).
- [3] Ulrich Kohlenbach. *Applied Proof Theory: Proof Interpretations and their Use in Mathematics. Proof interpretations and their use in mathematics*. Springer monographs in mathematics. Includes bibliographical references (p. [507]-523) and index. Berlin: Springer Berlin Heidelberg, 2008. 532 pp. ISBN: 9783540775331. DOI: [10.1007/978-3-540-77533-1](https://doi.org/10.1007/978-3-540-77533-1).
- [4] Eugene Lukacs. *Stochastic Convergence*. Ed. by E. Lukacs Z. W. Birnbaum. 2nd ed. Description based upon print version of record. Burlington: Elsevier Science, 1975. 215 pp. ISBN: 9780124598607. DOI: [10.1016/c2013-0-07416-6](https://doi.org/10.1016/c2013-0-07416-6).
- [5] Morenikeji Neri and Nicholas Pischke. *Proof mining and probability theory*. 2024. DOI: [10.48550/ARXIV.2403.00659](https://doi.org/10.48550/ARXIV.2403.00659).
- [6] Morenikeji Neri and Thomas Powell. “On quantitative convergence for stochastic processes: Crossings, fluctuations and martingales”. In: (2024). DOI: [10.48550/ARXIV.2406.19979](https://doi.org/10.48550/ARXIV.2406.19979).
- [7] Thomas Powell and Alex Wan. *An approximate zero-one law via the Dialectica interpretation*. 2025. DOI: [10.48550/ARXIV.2508.20849](https://doi.org/10.48550/ARXIV.2508.20849).